

Probability current

Let us consider the time derivative of particle density $\rho(x,t) = |\psi(x,t)|^2$

$$\frac{\partial \rho}{\partial t} = \frac{\partial}{\partial t} \psi^* \psi = \frac{\partial \psi^*}{\partial t} \psi + \psi^* \frac{\partial \psi}{\partial t}$$

Quantities $\frac{\partial \psi}{\partial t}$ and $\frac{\partial \psi^*}{\partial t}$ can be taken from the time-dependent Schrödinger equation and its complex conjugate:

$$\frac{\partial \psi}{\partial t} = \frac{1}{i\hbar} \left(-\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + V\psi \right)$$

$$\frac{\partial \psi^*}{\partial t} = \frac{1}{-i\hbar} \left(-\frac{\hbar^2}{2m} \frac{\partial^2 \psi^*}{\partial x^2} + V\psi^* \right)$$

When we substitute these into the expression for $\frac{\partial \rho}{\partial t}$ the terms containing V will cancel out (we assume $V^* = V$) and we get

$$\frac{\partial \rho}{\partial t} = \frac{-i\hbar}{2m} \left(\psi \frac{\partial^2 \psi^*}{\partial x^2} - \psi^* \frac{\partial^2 \psi}{\partial x^2} \right)$$

We can rewrite it as

$$\frac{\partial \rho}{\partial t} = -\frac{i\hbar}{2m} \frac{\partial}{\partial x} \left(\psi \frac{\partial \psi^*}{\partial x} - \psi^* \frac{\partial \psi}{\partial x} \right)$$

or

$$\frac{\partial \rho}{\partial t} = -\frac{\partial}{\partial x} j$$

where

$$j = \frac{i\hbar}{2m} \left(\psi \frac{\partial \psi^*}{\partial x} - \psi^* \frac{\partial \psi}{\partial x} \right)$$

$$= \frac{\hbar}{m} \text{Im} \left[\psi^* \frac{\partial \psi}{\partial x} \right]$$

$$= \frac{\hbar}{m} \text{Re} \left[\psi^* \hat{p} \psi \right]$$

If we did the same exercise in 3D by replacing

$$\frac{\partial^2 \psi}{\partial x^2} \rightarrow \nabla^2 \psi$$

we would have obtained

$$\begin{aligned} \frac{\partial \rho}{\partial t} &= -\nabla \cdot \vec{j} \quad \text{where} \quad \vec{j} = \frac{i\hbar}{2m} (\psi \nabla \psi^* - \psi^* \nabla \psi) \\ &= \frac{\hbar}{m} \text{Im} [\psi^* \nabla \psi] \\ &= \frac{\hbar}{m} \text{Re} [\psi^* \hat{p} \psi] \end{aligned}$$

The above equations look just like the familiar continuity equation where j (or \vec{j}) is a flux. This flux can be associated with the "probability current".

For charged particles (e.g. electron) the probability density is proportional to the charge density.

$$\rho_{\text{charge}} = q \rho_{\text{prob.}}$$

\swarrow
charge

Similarly the probability current is pretty much the same (up to a constant factor q) as the charge current.